Public Health Key Documents February 2017

NIHR

Autologous chondrocyte implantation (ACI) in the knee: systematic review and economic evaluation

The evidence base for ACI has improved since the last appraisal by NICE. In most analyses, the incremental cost-effectiveness ratios for ACI compared with microfracture appear to be within a range usually considered acceptable. Health Technology Assessment Systematic search: Yes February 2017 https://www.journalslibrary.nihr.ac.uk/hta/hta21060/#/abstract

KINGS FUND

Nil

HIS

Nil

SGHD

Demand Optimisation in Diagnostics

It has been recognised for many years that there is considerable variation in the use of diagnostic tests across the NHS. While some of this variation can be explained, there still exists considerable levels of inappropriate requesting by clinicians. Demand Optimisation can be applied to many diagnostic services, however the main focus of the conclusions and recommendations of this report relate to laboratory medicine. Scottish Government report Systematic search: No February 2017 http://www.gov.scot/Publications/2017/02/5322

SIGN

SIGN146 Cutaneous melanoma

This guideline has been revised to take account of new evidence in diagnostic and prognostic indicators, imaging techniques and laboratory investigation and systemic therapy. This guideline provides advice at all stages of the patient's pathway of care, from primary prevention to early recognition, treatment and follow up. It does not address melanomas of non-cutaneous origin such as melanomas arising from mucosae, ocular melanomas and other rare non-cutaneous sites. SIGN Guideline

HEALTH SCOTLAND

Nil

NICE

NG65 Spondyloarthritis in over 16s: diagnosis and management

This guideline covers diagnosing and managing spondyloarthritis that is suspected or confirmed in adults who are 16 years or older. It aims to raise awareness of the features of spondyloarthritis and provide clear advice on what action to take when people with signs and symptoms first present in healthcare settings. It also provides advice on the range of treatments available.

NICE Guidance Systematic search: Yes February 2017 https://www.nice.org.uk/guidance/ng65

NG64 Drug misuse prevention: targeted interventions

This guideline covers targeted interventions to prevent misuse of drugs, including illegal drugs, 'legal highs' and prescription-only medicines. It aims to prevent or delay harmful use of drugs in children, young people and adults who are most likely to start using drugs or who are already experimenting or using drugs occasionally. NICE Guidance Systematic search: Yes

February 2017 https://www.nice.org.uk/guidance/ng64

TA432 Everolimus for advanced renal cell carcinoma after previous treatment

Everolimus is recommended as an option for treating advanced renal cell carcinoma that has progressed during or after treatment with vascular endothelial growth factor targeted therapy.

NICE Technology Appraisal Systematic search: Yes February 2017 https://www.nice.org.uk/guidance/ta432

TA433 Apremilast for treating active psoriatic arthritis

Apremilast, alone or in combination with disease-modifying antirheumatic drugs (DMARDs), is recommended as an option for treating active psoriatic arthritis in adults only if they have peripheral arthritis with 3 or more tender joints and 3 or more swollen joints and their disease has not responded to adequate trials of at least 2 standard DMARDs, given either alone or in combination. NICE Technology Appraisal Systematic search: Yes February 2017 https://www.nice.org.uk/guidance/ta433

CG173 Neuropathic pain in adults: pharmacological management in nonspecialist settings

This guideline covers managing neuropathic pain with pharmacological treatments in adults in non-specialist settings. It aims to improve quality of life for people with conditions such as neuralgia, shingles and diabetic neuropathy by reducing pain and promoting increased participation in all aspects of daily living. The guideline sets out how drug treatments for neuropathic pain differ from traditional pain management NICE Clinical Guidance

Systematic search: Yes February 2017 https://www.nice.org.uk/guidance/cg173

CG190 Intrapartum care for healthy women and babies

This guideline covers the care of healthy women and their babies, during labour and immediately after the birth. It focuses on women who give birth between 37 and 42 weeks of pregnancy. The guideline helps women to make an informed choice about where to have their baby. It also aims to reduce variation in areas of care such as fetal monitoring during labour and management of the third stage of labour. NICE Clinical Guidance Systematic search: Yes February 2017 https://www.nice.org.uk/guidance/cg190

CG74 Surgical site infections: prevention and treatment

This guideline covers preventing and treating surgical site infections in adults, young people and children who are having a surgical procedure involving a cut through the skin. It recommends effective methods to use before, during and after surgery to minimise the risk of infection. NICE Clinical Guidance Systematic search: Yes

February 2017 https://www.nice.org.uk/guidance/cg74

CG139 Healthcare-associated infections: prevention and control in primary and community care

This guideline covers preventing and controlling healthcare-associated infections in children, young people and adults in primary and community care settings. It provides a blueprint for the infection prevention and control precautions that should be applied by everyone involved in delivering NHS care and treatment.

NICE Clinical Guidance Systematic search: Yes February 2017 https://www.nice.org.uk/guidance/cg139

CG146 Osteoporosis: assessing the risk of fragility fracture

This guideline covers assessing the risk of fragility fracture in people aged 18 and over with osteoporosis. It aims to provide guidance on the selection and use of risk assessment tools in the care of adults at risk of fragility fractures in all NHS settings. NICE Clinical Guidance Systematic search: Yes February 2017 https://www.nice.org.uk/guidance/cg146

DG27 Molecular testing strategies for Lynch syndrome in people with colorectal cancer

Offer testing to all people with colorectal cancer, when first diagnosed, using immunohistochemistry for mismatch repair proteins or microsatellite instability testing to identify tumours with deficient DNA mismatch repair, and to guide further sequential testing for Lynch syndrome. Do not wait for the results before starting treatment.

NICE Diagnostic Guidance Systematic search: Yes February 2017 https://www.nice.org.uk/guidance/dg27

ES5 Severe sialorrhoea (drooling) in children and young people with chronic neurological disorders: oral glycopyrronium bromide

Sialanar is licensed for short-term intermittent use and is only licensed in children. In trials, participants treated with glycopyrronium bromide had statistically significantly improved drooling after 8 weeks compared with placebo. NICE Evidence Summary Systematic search: Yes February 2017 https://www.nice.org.uk/advice/es5/chapter/Key-points

ES6 Parkinson's disease with motor fluctuations: safinamide

The main clinical benefits of safinamide at 24 weeks were an increase in 'on time' without troublesome dyskinesia, and a similar reduction in 'off time', compared with placebo. This effect was still observed at a 2-year follow-up. There are no studies comparing the efficacy and safety of safinamide with other active treatments, including other MAOB inhibitors. The NICE guideline on Parkinson's disease makes recommendations on the place in therapy of adjuvant treatments. NICE Evidence Summary Systematic search: Yes

February 2017 https://www.nice.org.uk/advice/es6/chapter/Key-points

EPPI Centre Nil

AHRQ (Agency for Healthcare Research and Quality - USA)

Health Foundation

Reducing hospital admissions by improving continuity of care in general practice

This briefing summarises research that analysed data from over 230,000 anonymised patient records for older people aged 62-82 years. Researchers found there were fewer hospital admissions for certain conditions when patients saw the same GP more consistently. Patients seeing their usual GP two more times out of every 10 was associated with 6% fewer avoidable hospital admissions. Health Foundation briefing

Systematic search: No February 2017 <u>http://www.health.org.uk/publication/reducing-hospital-admissions-improving-</u> <u>continuity-care-general-practice</u>

Canadian Agency for Drugs and Technologies in Health (CADTH)

Blue Light Cystoscopy in Patients with Suspected Non-Muscle Invasive Bladder Carcinoma: A Review of Clinical Utility

Hexaminolevulinate-guided transurethral resection of bladder tumors was associated with a decreased risk of bladder cancer recurrence and a decreased risk of progression to muscle invasive bladder cancer compared to white light cystoscopy.

CADTH Rapid Response Report Systematic search: Limited February 2017 <u>https://www.cadth.ca/blue-light-cystoscopy-patients-suspected-non-muscle-invasive-</u> bladder-carcinoma-review-clinical

The Use of Osteochondral Allograft for the Ankle, Knee, and Shoulder: Clinical Effectiveness and Cost-Effectiveness

For all three joints, while the operation can improve pain and functional outcomes, with a high level of patient satisfaction, high rates of complications, reoperations, and clinical failures were reported.

CADTH Rapid Response Report Systematic search: Limited February 2017

https://www.cadth.ca/use-osteochondral-allograft-ankle-knee-and-shoulder-clinicaleffectiveness-and-cost-effectiveness

Aripiprazole for Borderline Personality Disorder: A Review of the Clinical Effectiveness

Limited evidence indicates that aripiprazole appears to be a safe and effective agent in the long term treatment of patients with borderline personality disorder. Similar clinical effectiveness and safety profiles were reported comparing aripiprazole with olanzapine in the treatment of BPD.

CADTH Rapid Response Report Systematic search: Limited February 2017 https://www.cadth.ca/aripiprazole-borderline-personality-disorder-review-clinicaleffectiveness-0

McGill University Health Centre (Canada)

Nil

Health Information & Quality Authority (Ireland)

Mechanical Thrombectomy for Stroke

In selected patients, mechanical thrombectomy using stent retriever devices is a safe and effective procedure when provided as an adjunct to standard medical care within six to 12 hours of onset of an acute ischaemic stroke. Health Technology Assessment Systematic search: Yes February 2017 https://www.hiqa.ie/reports-and-publications/health-technology-assessments/htamechanical-thrombectomy-stroke

Campbell Collaboration

12-step programs for reducing illicit drug use

The results of this review suggest that 12-step interventions to support illicit drug users are as effective as alternative psychosocial interventions in reducing drug use. However, the evidence is limited, estimated effect sizes are small, and other influencing factors may not have been totally considered.. Systematic Review Systematic search: Yes February 2017 https://www.campbellcollaboration.org/library/12-step-programmes-illicit-drugabuse-reduction.html

Glasgow Centre for Population Health

Nil

NICE FORWARD PLANNING – Publications due March 2017

Hepatocellular carcinoma (advanced and metastatic) - sorafenib (first line) Single Technology Appraisal

Breast cancer (refractory, HER2 positive) - trastuzumab-emtansine Single Technology Appraisal

Head and neck cancer (squamous cell carcinoma) - cetuximab

Single Technology Appraisal

Mental health of adults in contact with the criminal justice system Clinical Guideline

Familial Breast Cancer Clinical Guideline

Managing medicines for adults receiving social care in the community Clinical Guideline

Waldenstrom's macroglobulinaemia - ibrutinib Single Technology Appraisal

ENDURALIFE-powered CRT-D devices for treating heart failure Medical Technology

Hypophosphatasia (paediatric-onset) - asfotase alfa Highly Specialised Technology Evaluation